

(A) One way ANOVA table:- (Single factor ANOVA)  
 It is convenient to summarize the result of an analysis of variance in a table.

Source of variation	Sum of Square (SS)	DOF	Mean Square MS	F-Ratio
Between samples	$SS_B$	$K-1$	$MS_B = SS_B / DOF_B$	$MS_B / MS_W$
Within samples	$SS_W$	$n-K$	$MS_W = SS_W / DOF_W$	
Total	$SS_T$	$n-1$		

(Qn)

$X_1$	$X_2$	$X_3$
1	2	2
2	4	3
5	2	4

Ans

Step-1:  $H_0: \mu_1 = \mu_2 = \mu_3$   
 $H_a: \mu_1 \neq \mu_2 \neq \mu_3$   
 $\alpha = 0.05$

Step-2: Dof between =  $(k-1)$   
 where,  $k = \text{No of groups}$

So,  $k = 3$  //  $3-1 = 2$  //

Dof within =  $(n-k)$ , where  $n = \text{No of scores}$   
 $n = 9$

So,  $(n-k) = (9-3) = 6$  //

Dof Total =  $(n-1)$  So,  $(9-1) = 8$  //

Table value (Refer the table)

Dof between  $\rightarrow$  Numerator

Dof within  $\rightarrow$  Denominator

So Table value = 5.14 //

Step-3:

$X_1$	$X_2$	$X_3$	$X$
1	2	2	1
2	4	3	2
5	2	4	5
<u>5</u>	<u>2</u>	<u>4</u>	2
$\sum X_1 = 8$	$\sum X_2 = 8$	$\sum X_3 = 9$	4
$\bar{X}_1 = 8/3$	$\bar{X}_2 = 8/3$	$\bar{X}_3 = 9/3$	2
$= 2.67$	$= 2.67$	$= 3$	2
			3
			4
			<u>4</u>
			$\sum X = 25$

~~Calculate~~  $G\bar{X} = 25/9$   
 $= 2.78$

Step-4: (1) Calculate Sum of Square Total (SST):-

$SST = \text{Sum Square } (x_i - G\bar{X}) \text{ of}$   
 $\text{Sum of Square Total } (x_i - G\bar{X})^2$



$x_1$	$x_2$	$x_3$	$SS_T$
$(1 - 2.78)^2$ $= 3.16$	0.60	0.60	$3.16 + 0.60 +$
$(2 - 2.78)^2$ $= 0.60$	1.49	0.05	$4.94 + 0.60 +$
$(5 - 2.78)^2$ $= 4.94$	0.60	1.49	$1.49 + 0.60 +$
			$0.60 + 0.05 +$
			1.49
			<u><u>13.56</u></u>

(II) Calculate Sum of Square within ( $SS_W$ ):-

$SS_W =$  Sum of Square  $(x_i - \bar{x}_i)$  of  
Sum of Square within  $(x_i - \bar{x}_i)^2$

$x_1$	$x_2$	$x_3$	$SS_W$
$(1 - 2.67)^2$ $= 2.78$	$(2 - 2.67)^2$ $= 0.44$	$(2 - 3)^2$ $= 1$	$2.78 + 0.44 +$
0.44	1.78	0	$5.44 + 0.44 +$
5.44	0.44	1	$1.78 + 0.44 +$
			$1 + 0 + 1$
			<u><u>13.33</u></u>

(III) Calculate Sum of Square Between ( $SS_B$ )

$$SS_B = SS_T - SS_W = 13.56 - 13.33 = \underline{\underline{0.22}}$$

Step - 5 :- Calculate Mean Variance

(I) Mean Square Between ( $MS_B$ )

$$\Rightarrow MS_B = SS_B / \text{Dof}_B = 0.22 / 2 = \underline{\underline{0.11}}$$

(II) Mean Square within ( $MS_W$ )

$$\begin{aligned} \Rightarrow MS_W &= SS_W / \text{Dof}_W \\ &= 13.33 / 6 = \underline{\underline{2.22}} \end{aligned}$$


Step-6 Calculate f-Ratio:-

$$f = MS_R / MS_W = 0.11 / 2.22 = \underline{\underline{0.05}}$$

Critical factor of f value from the table = 5.14

Conclusion:-

Calculated value  $<$  Table value



$$0.05 < 5.14$$

accepted

Hence  $H_0$  fails to reject i.e.,  $H_0$  is accepted.  
So there is no significant difference between all the variables of three different series.

---



(B) Two way ANOVA table:-

(Two factor analysis)

It takes the following form.

Source of Variance	Sum of Square (SS)	Dof	Mean Square (MS)	F Ratio
Between Rows	$SS_R$	$(r-1)$	$MS_R = \frac{SS_R}{Dof_r}$	$\frac{MS_R}{MS_E}$
Between Columns	$SS_c$	$(c-1)$	$MS_c = \frac{SS_c}{Dof_c}$	$\frac{MS_c}{MS_E}$
Error Residuals	$SS_E$	$(r-1) \times (c-1)$	$MS_E = \frac{SS_E}{Dof_E}$	
Total	$SS_T$	$(rc-1)$		

On

fertilizers ↓	seeds →		
	a	b	c
w	6	5	5
x	7	5	4
y	3	3	3
z	8	7	4



Step-1 Calculate Correction factor:-

$$\text{Correction factor} = T^2 / n$$

where,  $T = \text{Sum of all values}$

$$(6+7+3+8+5+5+3+7+5+4+3+4) = 60$$

$$n = R \times C = 4 \times 3 = 12$$

$$\text{Correction factor} = (60)^2 / 12 = 300 //$$

Step-2 Calculate Sum of Square Total (SST):-

$$SST = (\text{Total Sum Square} - \text{Correction factor})$$

w	36	25	25
x	49	25	16
y	9	9	9
z	64	49	16

$$\begin{aligned} \text{Total Sum Square} \\ = 332 \end{aligned}$$

$$SST = 332 - 300 = 32 //$$

Step-3 Calculate SSc, SSR & SSe

	R	C	T	CR	CCR = SSR / ni
w	6	5	5	16	$16^2 / 3 = 85.33$
x	7	5	4	16	$16^2 / 3 = 85.33$
y	3	3	3	9	$9^2 / 3 = 27$
z	8	7	4	19	$19^2 / 3 = 120.33$
Sc	24	20	16		SSR = 317.66
SSc = $\frac{Sc^2}{ni}$	$\frac{24^2}{4} = 144$	$\frac{20^2}{4} = 100$	$\frac{16^2}{4} = 64$	SSe = 308	



Step - 4

(i) Calculate SS between Columns treatment:-  
 $= SS_C - \text{Correction factor}$   
 $= 308 - 300 = 8 //$

(ii) Calculate SS between Row treatment:-  
 $= SS_R - \text{Correction factor}$   
 $= 317.66 - 300 = 17.66 = 18 //$

(iii) Calculate SS residual & Error:-  
 $= SS_T - (SS_C + SS_R)$   
 $= 32 - (8 + 18) = 6 //$

Step - 5

ANOVA Table

Source of Variation	SS	Dof	MS	F-Ratio
Between Row	18	$(4-1) = 3$	$18/3 = 6$	$6/1 = 6$
Between Column	8	$(3-1) = 2$	$8/2 = 4$	$4/1 = 4$
Residual Error	6	$3 \times 2 = 6$	$6/6 = 1$	//////
Total	32	$(3 \times 2) - 1 = 5$	//////	//////

N.B:- Critical value (Ref to table)

D.O.F of (Row, Residual Error) are numerator & denominator respectively as well as (Column, Residual Error) are also numerator & denominator respectively. ie  $F(2, 6) = 5.14 //$   
&  $F(3, 6) = 4.76 //$



Decision:-

Column:-

Hence  $f$ -value calculated in terms of between Column  $<$  critical value of Column  
 $\Rightarrow 4 < 5.14$

Hence  $H_0$  fails to reject ( $H_0$  accepted)

So there is no significant difference between seeds.


Row:-

Hence  $f$  value calculated in terms of row  $>$  critical value of row  
 $\Rightarrow 6 > 4.76$

Hence  $H_0$  is rejected


So there is significant difference between the fertilizers

(seeds)



Accepting  
 $H_0$

(fertilizer)



Rejected  
 $H_0$



**The F - Distribution with  $\alpha = 0.05$**

$\nu_2 \setminus \nu_1$	2	3	4	5	6	7	8	9
2	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38
3	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
4	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
5	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
6	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
7	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68
8	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39
9	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18